arXiv:math/0501538v1 [math.AC] 29 Jan 2005

On the generating poset of Schubert cycles and the characterization of Gorenstein property

Mitsuhiro MIYAZAKI[1]

Abstract: The homogeneous coordinate ring of a Schubert variety (a Schubert cycle for short) is an algebra with straightening law generated by a distributive lattice. This paper gives a simple method to study the set of all the join-irreducible elements of this distributive lattice, and gives a simple proof of the criterion of the Gorenstein property of Schubert cycles.

Key words: Schubert cycle, ASL, distributive lattice, Gorenstein property

1 Introduction

In this note, all rings and algebras are assumed to be commutative with identity element. Let m and n be integers with $1 \leq m \leq n$, B a ring and S a B-algebra. For an $m \times n$ matrix M with entries in S, we denote by $G(M)$ the B-subalgebra of S generated by all the maximal minors of M.

From now on, we assume that B is a field. Let X be the $m \times n$ matrix of indeterminates. Then it is well known that $G(X)$ is the homogeneous coordinate ring of the Grassmann variety $G_{m,n}$ of m-planes in n-space ([HP], [DEP], [BV]). And it is also well-known that $G(X)$ is a graded algebra with straightening law (graded ASL for short) over B generated by the poset $\Gamma(X)$, where

$$\Gamma(X) := \{[a_1, \ldots, a_m] \mid 1 \leq a_1 < \cdots < a_m \leq n\}$$

([DEP], [BV]). The order of $\Gamma(X)$ is defined by

$$[a_1, \ldots, a_m] \leq [b_1, \ldots, b_m] \overset{\text{def}}{\Longleftrightarrow} a_i \leq b_i \text{ for } i = 1, \ldots, m.$$

And the element $[a_1, \ldots, a_m]$ of $\Gamma(X)$ is mapped to $\det(X_{ia_j}) \in G(X)$. It is easily verified that $\Gamma(X)$ is a distributive lattice.

Now we fix an n-dimensional vector space V and a complete flag $V = V_n \supsetneq V_{n-1} \supsetneq \cdots \supsetneq V_1 \supsetneq V_0 = 0$ of subspaces. For integers $a_1, \ldots, a_m$ with $1 \leq a_1 < \cdots < a_m \leq n$, the Schubert subvariety $\Omega(a_1, \ldots, a_m)$ of $G_{m,n}$ is defined by

$$\Omega(a_1, \ldots, a_m) := \{W \in G_{m,n} \mid \dim(W \cap V_{a_i}) \geq i \text{ for } i = 1, \ldots, m\}.$$

---

[1] Dept. Math. Kyoto University of Education, 1 Fukakusa-Fujinomori-cho, Fushimi-ku, Kyoto, 612-8522 Japan, E-mail:`g53448@kyokyo-u.ac.jp`

If we put $b_i = n + 1 - a_{m+1-i}$ for $i = 1, \dots, m$, $\gamma = [b_1, \dots, b_m]$ and $\Gamma(X;\gamma) = \{\delta \in \Gamma(X) \mid \delta \geq \gamma\}$, then the homogeneous coordinate ring of the Schubert variety (Schubert cycle for short) is

$$G(X;\gamma) := G(X)/(\Gamma(X) \setminus \Gamma(X;\gamma))G(X).$$

This ring is a graded ASL over B generated by $\Gamma(X;\gamma)$ ([DEP], [BV]).

Set

$$U_\gamma := \begin{pmatrix} 0 & \cdots & 0 & U_{1b_1} & \cdots & U_{1b_2-1} & U_{1b_2} & \cdots & \cdots & \cdots & \cdots & U_{1n} \\ 0 & \cdots & 0 & 0 & \cdots & 0 & U_{2b_2} & \cdots & \cdots & \cdots & \cdots & U_{2n} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \\ 0 & \cdots & 0 & 0 & \cdots & 0 & 0 & \cdots & 0 & U_{mb_m} & \cdots & U_{mn} \end{pmatrix},$$

where U_{ij} are independent indeterminates. Then by [BV, (7.16)], $G(X;\gamma) \simeq G(U_\gamma)$. So a Schubert cycle is a graded ASL generated by a distributive lattice and an integral domain.

Now we recall the following fact.

Fact 1.1 ([Sta], [Hib]) *Let A be a graded ASL domain over a field generated by a distributive lattice D. Assume that* $\deg(\alpha) + \deg(\beta) = \deg(\alpha \wedge \beta) + \deg(\alpha \vee \beta)$ *for all α, $\beta \in D$. Then A is Gorenstein if and only if the set of all the join-irreducible elements of D is pure.*

Recall that an element x in a lattice L is said to be join-irreducible if there is unique y in L such that $y \lessdot x$, where $y \lessdot x$ means that $y < x$ and there is no element $z \in L$ such that $y < z < x$.

So in this note, we shall study the set P of all the join-irreducible elements of $\Gamma(X;\gamma)$ and give a simple proof of the criterion of the Gorenstein property of Schubert cycles.

2 Join-irreducible elements of $\Gamma(X;\gamma)$

As in the previous section, let m and n be integers such that $1 \leq m \leq n$ and $\gamma = [b_1, \dots, b_m] \in \Gamma(X)$. And let P be the set of all the join-irreducible elements of $\Gamma(X;\gamma)$.

Assume that $[c_1, \dots, c_m]$, $[d_1, \dots, d_m] \in \Gamma(X)$. Then it is easy to verify that

$$[c_1, \dots, c_m] \lessdot [d_1, \dots, d_m] \iff \exists i; d_i = c_i + 1, d_j = c_j (j \neq i).$$

So $[c_1, \dots, c_m]$ is a join-irreducible element of $\Gamma(X;\gamma)$ if and only if there is a unique i such that

$$c_i > b_i, \quad c_i > c_{i-1} + 1, \tag{1}$$

where we set $c_0 := 0$. For a join-irreducible element $[c_1, \dots, c_m]$ of $\Gamma(X;\gamma)$, we take i satisfying (1), and set

$$p := n - c_i - (m - i), \quad q := i - 1.$$

That is, $p = \#\{t \in \{1, 2, \dots, n\} \mid t > c_i \text{ and } t \notin \{c_1, c_2, \dots, c_m\}\}$ and $q = \#\{t \in \{1, 2, \dots, n\} \mid t < c_i \text{ and } t \in \{c_1, c_2, \dots, c_m\}\}$. We denote the map which send $[c_1, \dots, c_m]$ to (p, q) by φ.

It is easy to construct a join-irreducible element $[c_1, \dots, c_m]$ such that $\varphi([c_1, \dots, c_m]) = (p, q)$, if (p, q) is in the image of φ. And it also easy to verify that if (p, q) is in the image of φ and $0 \le p' \le p$, $0 \le q' \le q$, then (p', q') is also in the image of φ. Moreover, if $[c_1, \dots, c_m]$ and $[d_1, \dots, d_m]$ are join-irreducible elements of $\Gamma(X;\gamma)$, and $\varphi([c_1, \dots, c_m]) = (p, q)$, $\varphi([d_1, \dots, d_m]) = (p', q')$, then

$$[c_1, \dots, c_m] \le [d_1, \dots, d_m] \iff p \ge p', q \ge q'.$$

In particular, P is isomorphic to a finite filter (i.e. a subset F of a poset such that $x \in F$ and $y > x$ implies $y \in F$) of $\boldsymbol{N} \times \boldsymbol{N}$ whose order is defined by

$$(p, q) \le (p', q') \overset{\text{def}}{\iff} p \ge p' \text{ and } q \ge q'.$$

And if $\varphi([c_1, \dots, c_m]) = (p, q)$, then the coheight of $[c_1, \dots, c_m]$ is $p + q$.

Therefore, if we set

$$\{i \mid b_i + 1 < b_{i+1},\ b_i < n\} = \{l_1, \dots, l_u\}, \quad l_1 < \cdots < l_u,$$

then the minimal elements of P are $[b_1, \dots, b_{l_1 - 1}, b_{l_1} + 1, b_{l_1 + 1}, \dots b_m]$, …, $[b_1, \dots, b_{l_u - 1}, b_{l_u} + 1, b_{l_u + 1}, \dots, b_m]$ and their coheights are $n - m - b_{l_1} + 2l_1 - 2$, …, $n - m - b_{l_u} + 2l_u - 2$ respectively.

Example 2.1 If $n = 14$, $m = 7$, $[b_1, \dots, b_m] = [2, 4, 5, 9, 10, 12, 13]$, then $u = 4$, $l_1 = 1$, $l_2 = 3$, $l_3 = 5$ and $l_3 = 7$ and the minimal elements of P are

$$\begin{aligned}\gamma_1 &= [3, 4, 5, 9, 10, 12, 13]\\ \gamma_2 &= [2, 4, 6, 9, 10, 12, 13]\\ \gamma_3 &= [2, 4, 5, 9, 11, 12, 13]\\ \gamma_4 &= [2, 4, 5, 9, 10, 12, 14].\end{aligned}$$

And the Hasse diagram of P is the following.



Note that this poset is the poset obtained by Stanley [Sta, 5.b].

Using Fact 1.1, we obtain a simple proof of the following criterion of Gorenstein property of Schubert cycles.

Theorem 2.2 *In the notation above, $G(X;\gamma)$ is Gorenstein if and only if $b_{l_i} - 2l_i$ is constant for $i = 1, \ldots, u$.*

References

[BV] Bruns, W. and Vetter, U.: "Determinantal Rings." Lecture Notes in Mathematics **1327** Springer (1988)

[DEP] DeConcini, C., Eisenbud, D. and Procesi, C.: "Hodge Algebras." Astérisque **91** (1982)

[Hib] Hibi, T.: *Distributive lattices, affine smigroup rings and algebras with straightening laws.* in "Commutative Algebra and Combinatorics" (M. Nagata and H. Matsumura, ed.), Advanced Studies in Pure Math. **11** North-Holland, Amsterdam (1987), 93–109.

[HP] Hodge, W. V. D. and Pedoe, D.: "Methods of algebraic geometry" Vol. 2 Cambridge University Press, Cambridge, 1968

[Sta] Stanley, R. P.: *Hilbert Functions of Graded Algebras.* Adv. Math. **28** (1978), 57–83.